
















Additional File 2: Full CLIP-Seq results










On the following pages, the motifs that were recovered by *MEMERIS*, *RNAcontext*, *GraphProt*, and ssHMM from each of the 25 CLIP-Seq datasets are shown. The table has the following six columns:









1. Protein - Name of the RBP (with protocol in parentheses if multiple CLIP-Seq datasets exist for the protein)
2. Literature motif - Motifs published by other studies (if present)
3. ssHMM - Visualization of the ssHMM trained on the CLIP-Seq dataset. Below the visualization, four numbers are printed: the sequence, structure, and combined information content for comparison with *GraphProt* (in red) and the sequence information content for comparison with *MEMERIS* and *RNAcontext* (in blue)
4. MEMERIS - Sequence motif by *MEMERIS*. Below, the information content of the sequence logo is printed.
5. RNAcontext - Sequence motif by *RNAcontext*. Below, the information content of the sequence logo is printed.
6. GraphProt - Sequence and structure motif by *GraphProt*. Below, the sequence, structure, and combined information content of the logos is printed.









Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
Ago1234					
		0.69/0.86/1.82/0.34	0.91	0.04	0.64/0.34/0.95
Ago2 (HITSCLIP)					
		0.67/0.85/1.76/0.42	0.91	0.15	0.56/0.31/0.84

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
Ago2 (PARCLIP)		1 2 3 4 5 6			
	Multiloop				
	Hairpin				
	Stem				
	Internal loop				
	Exterior loop				
		0.88/0.9/1.87/0.52	0.92	0.21	0.8/0.28/1.04
CAPRIN1		1 2 3 4 5 6			
	Multiloop				
	Hairpin				
	Stem				
	Internal loop				
	Exterior loop				
		0.7/0.84/1.82/0.24	0.93	0.01	0.47/0.57/1.03









Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
DGCR8					
		1.4/1.2/2.71/0.66	1.04	1.06	0.76/1.49/2.2
DICER			runtime exceeded 7 days		
		0.91/1.62/2.8/0.31		0.21	0.65/1.75/2.38










Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
EIF4A3		 <p>1.45/0.99/2.56/0.74</p>	 <p>1.12</p>	 <p>0.27</p>	 <p>1.35/0.58/1.89</p>
EWSR1	AU-rich stem-loops [7]	 <p>1.07/1.11/2.25/0.49</p>	 <p>1.03</p>	 <p>0.18</p>	 <p>0.52/0.37/0.91</p>











Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
EZH2					
					
					
					
		0.84/1.66/2.55/0.55	0.93	0.27	0.74/1.37/2.05
FMRP					
					
					
					
		0.98/1.83/2.84/0.67	0.97	0.47	0.8/1.53/2.32









Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
FXR2					
		0.88/1.48/2.42/0.4	0.88	0.63	0.71/0.62/1.29
HuR (HITSCLIP)	AU-rich regions [1]				
		2/1.33/3.32/1.05	1.45	0.28	1.74/0.77/2.41









Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
HuR (PARCLIP)	AU-rich regions [1]				
		2/0.91/2.9/1.01	1.39	0.37	1.79/0.36/2.06
IGF2BP1/2/3	 [2]				
		1.14/0.85/2.08/0.7	1.08	0.64	0.87/0.32/1.19





Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
LIN28B					
		1.03/1.3/2.37/0.49	0.96	0.33	0.68/0.42/1.09
MOV10					
		1.03/0.77/2.17/0.42	1.00	0.01	0.86/1.33/2.14

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
Nova	UCAUY [3]				
		1.7/1.04/2.75/1.05	1.31	0.46	0.77/0.44/1.14
PTBP1	 [4]				
		1.54/1.44/2.99/0.82	1.28	0.24	1.69/0.43/2.02

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
PUM2	 [2]	 1.54/1.26/2.89/1.07	 WebLogo 3.5.0	 WebLogo 3.4	 1.11/1.68/2.7
QKI	 [2]	 1.56/1.17/2.6/1.02	 WebLogo 3.5.0	 WebLogo 3.4	 0.73/0.52/1.22

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
SRSF1/SF2	 [6]	 1.44/1.59/3.42/0.74	runtime exceeded 7 days	 0.07	 1.48/0.29/1.79
TAF2N	AU-rich stem-loops [7]	 1.12/1.59/2.76/0.5	 1.07	 0.65	 0.59/0.38/0.99

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
TIA1	U-rich region [8]				
		1.92/0.68/2.61/0.94	1.38	0.47	1.72/0.37/2
YY1					
		0.84/1.33/2.84/0.52	1.21	0.23	1.09/1.6/2.61

Protein	Literature motif	ssHMM	MEMERIS	RNAcontext	GraphProt
ZC3H7B					
		0.93/0.78/1.97/0.41	1.03	0.33	0.64/0.68/1.28

Bibliography

- [1] Peng, S.S.-Y., Chen, C.-Y.A., Xu, N., Shyu, A.-B.: Rna stabilization by the au-rich element binding protein, hur, an elav protein. *The EMBO journal* **17**(12), 3461–3470 (1998)
- [2] Hafner, M., Landthaler, M., Burger, L., Khorshid, M., Hausser, J., Berninger, P., Rothballer, A., Ascano, M., Jungkamp, A.-C., Munschauer, M., Ulrich, A., Wardle, G.S., Dewell, S., Zavolan, M., Tuschl, T.: Transcriptome-wide Identification of RNA-Binding Protein and MicroRNA Target Sites by PAR-CLIP. *Cell* **141**(1), 129–141 (2010)
- [3] Jensen, K.B., Dredge, B.K., Stefani, G., Zhong, R., Buckanovich, R.J., Okano, H.J., Yang, Y.Y.L., Darnell, R.B.: Nova-1 regulates neuron-specific alternative splicing and is essential for neuronal viability. *Neuron* **25**(2), 359–371 (2000)
- [4] I, P., Ch, L., Jg, M., Jg, P.: Mutation of PTB binding sites causes misregulation of alternative 3' splice site selection in vivo., Mutation of PTB binding sites causes misregulation of alternative 3' splice site selection in vivo. *RNA* **3** 3(7, 7), 764–764778 (1997)
- [5] Xue, Y., Zhou, Y., Wu, T., Zhu, T., Ji, X., Kwon, Y.-S., Zhang, C., Yeo, G., Black, D.L., Sun, H., Fu, X.-D., Zhang, Y.: Genome-wide Analysis of PTB-RNA Interactions Reveals a Strategy Used by the General Splicing Repressor to Modulate Exon Inclusion or Skipping. *Molecular Cell* **36**(6), 996–1006 (2009)
- [6] Sanford, J.R., Wang, X., Mort, M., VanDuyn, N., Cooper, D.N., Mooney, S.D., Edenberg, H.J., Liu, Y.: Splicing factor SFRS1 recognizes a functionally diverse landscape of RNA transcripts. *Genome Research* **19**(3), 381–394 (2009)
- [7] Hoell, J.I., Larsson, E., Runge, S., Nusbaum, J.D., Duggimpudi, S., Farazi, T.A., Hafner, M., Borkhardt, A., Sander, C., Tuschl, T.: RNA targets of wild-type and mutant FET family proteins. *Nature Structural & Molecular Biology* **18**(12), 1428–1431 (2011)
- [8] Dember, L.M., Kim, N.D., Liu, K.-Q., Anderson, P.: Individual RNA Recognition Motifs of TIA-1 and TIAR Have Different RNA Binding Specificities. *Journal of Biological Chemistry* **271**(5), 2783–2788 (1996)