

Supplementary Table ST.8. Urinary hippurate concentration distribution over enterotypes (N=271). The statistical difference over the four enterotypes was tested using a non-parametric Kruskal-Wallis test, followed by post hoc Dunn tests for pairwise differences (two-sided testing).

X	Y	Kruskal-Wallis			Score	Mean	Std Err	Effect					
		ChiSquar	e	DF				Level	Level2	Difference	Dif	Z	size (r)
Enterotypes	Hippurate	41.7878		3	4.45E-09	Rum	Bact2	92.14	16.69	5.52	0.45524	2.04E-07	147
Enterotypes	Hippurate	41.7878		3	4.45E-09	Prev	Bact2	83.38	18.36	4.54	0.4984	3.36E-05	83
Enterotypes	Hippurate	41.7878		3	4.45E-09	Rum	Bact1	50.86	11.9	4.28	0.31183	0.00011	188
Enterotypes	Hippurate	41.7878		3	4.45E-09	Prev	Bact1	42.1	14.14	2.98	0.26733	0.01747	124
Enterotypes	Hippurate	41.7878		3	4.45E-09	Rum	Prev	8.75	12.68	0.69	0.05197	1	176
Enterotypes	Hippurate	41.7878		3	4.45E-09	Bact2	Bact1	-41.26	17.83	-2.31	-0.23747	0.12383	95