

# Clonal expansion and epigenetic inheritance of long-lasting NK cell memory

In the format provided by the  
authors and unedited

**Supplementary Table 1 Healthy blood donor characteristics.** HCMV serostatus was determined by anti-HCMV IgG ELISA (see Methods).

Donor ID	HCMV Serostatus	Sex	Age at inclusion	scATAC	scRNA
P1	+	m	45	x	x
P2	+	m	27	x	x
P3	+	w	29	x	x
P4	+	w	27	x	x
P5	+	m	28		x
N1	-	w	30	x	x
N2	-	m	29	x	x
N3	-	m	33	x	
N4	-	w	29	x	

**Supplementary Table 2 KIR ligand genotyping results.** KIR ligand genotypes were determined by sequence-specific priming PCR (see Methods).

Donor ID	HLA-A	HLA-B	HLA-C	Educating KIRs
P1	Bw4 <sup>-</sup>	Bw4 <sup>+</sup> Ile80	C1/C2	KIR2DL1, KIR2DL2/3, KIR3DL1
P2	Bw4 <sup>+</sup>	Bw4 <sup>+</sup> Ile80	C2/C2	KIR2DL1, KIR3DL1
P3	Bw4 <sup>+</sup>	Bw4 <sup>+</sup> Thr80	C1/C2	KIR2DL1, KIR2DL2/3, KIR3DL1
P4	Bw4 <sup>-</sup>	Bw4 <sup>-</sup>	C1/C2	KIR2DL1, KIR2DL2/3